

A New Formula for Depigmenting Human Skin

Albert M. Kligman, MD, PhD, Isaac Willis, MD

**Complete depigmentation of the normal
skin of adult male blacks was procured by**

to pigmentary abnormalities as "cos-
metic." One untoward effect of such

period began when Oettel produced
grey hair in black cats by feeding

Fig 1.—Left, Transverse section of skin before treatment. Right, Specimen from depigmented skin after six weeks of treatment. Epidermis is acanthotic and keratinocytes larger; granular layer is increased, with conspicuous granules (hematoxylin-eosin, $\times 630$).

of leukoderma at distant sites is a puzzling but rather characteristic adverse happening with this class of compounds.⁸

Moreover, MBEH proved to be a potent contact sensitizer. Worst of all, about half the subjects had leukoderma for the next two years and about a third for at least four years, presumably the longest period of follow-up. Persistent and perhaps permanent depigmentation is the spectre that caused the editorial comment that "the therapeutic use of MBEH has been responsible for a number of cosmetic disasters."⁹

Meanwhile, investigators were frenetically searching for chemicals that might be both safe and effective when they were applied topically. Brun evaluated the effect of thirty compounds applied on the pigmented nipples of guinea pigs and found that the monomethyl and monomethyl ethers of hydroquinone also were potent depigmenting agents.¹⁰ Chavin and Schlesinger injected a number of chemicals into goldfish and added certain mercaptoamines to the list of powerful depigmentors.² The latter are also effective in the treatment of mammalian skin; however, they are both irritating and malodorous.¹¹



Bleehen et al applied all the aforementioned compounds and many related ones, thirty-three in all, to black guinea pigs and discovered 4-isopropyl catechol to be the most effective.¹²

Fig 2.—Left, Before treatment. Melanin granules are prominent in basal layer, often in form of caps over the nuclei. Pigment granules are visible in horny layer of blacks. Right, After eight weeks of treatment. Pigment granules, mainly in basal cells, can barely be seen (Fontana's silver stain, $\times 630$).

Therapeutic Effectiveness of Basic Formula

Disorder	No. of Patients Treated	No. of Patients With Satisfactory Response
Melasma	16	14
Ephelides	11	
Postinflammatory pigmentation	18	12
Senile lentigines (hands)		

This, too, is a moderately strong contact allergen.



notes, is not a substituted phenol. Earlier, Denton et al were able to show that most of these compounds could block the hydroxylation of tyrosine to dopa in vitro and thus might inhibit the first step in melanin synthesis.³

Spencer noted that hydroquinone

sistent hyperpigmentation. Extensive patch testing of black volunteers with various irritating and allergenic chemicals awakened our sensibilities to the problem of hyperpigmentation in blacks.

Two observations led us to suppose that a way might yet be found to en-

chemical destination. It would serve no purpose to chronicle our experimental wanderings. The formula that eventuated has the following composition: tretinoin, 0.1%; hydroquinone, 5.0%; and dexamethasone, 0.1%.

The vehicle was either hydrophilic ointment or a solution consisting of equal parts

itself was apparently free of the deleterious effects of MBEH. With the use of 2%, 3%, and 5% concentrations of hydroquinone, depigmentation of senile lentigines in more than 75% of white males was obtained.¹⁸ Repigmentation occurred promptly after cessation of treatment and depigmentation of distant sites was absent. As shown by light microscopy, melanin granules were decreased in depigmented skin but the number of melanocytes was about the same. In this way the use of hydroquinone in a low concentration was introduced into practice.

In a study of 56 patients with various pigmentary disturbances, Arndt and Fitzpatrick found 2% and 5% hy-

hance the pharmacologic effectiveness of hydroquinone. First, in treating hundreds of acne patients with topically applied tretinoin, we became aware that the skin sometimes became lighter after many months of use. The degree of hypopigmentation was slight. Second, the intradermal injection of corticosteroids into the skin of blacks in connection with assays of anti-inflammatory activity quite commonly resulted in striking loss of pigment at the injection site. Cahn noticed this phenomenon after injection of corticosteroids into lesions of black patients.²⁰ We too have observed this, so far only in blacks. In time the pigment returns. The application of tape containing cortico-

were considerably less effective included petrolatum, lanolin, and polyethylene glycol. Dexamethasone was selected as the test steroid because it was economically available to us in a pure form. Other steroids may be substituted.

The cream and solution were never more than 30 days old. The addition of anti-oxidants did not enhance activity.

Method of Application

The agents were applied twice daily to squares of back skin outlined by a cardboard stencil. We came to appreciate that a moderate degree of irritation (peeling and erythema) promoted earlier and greater depigmentation. The amount applied was therefore regulated to achieve this effect.

Unpublished Studies

mentation.

Peeling and sometimes redness usually preceded depigmentation. Tenderness and burning were frequent complaints but inflammation was never severe enough to discontinue treatment. After about one month of treatment, the skin became hardened; thereafter, liberal applications produced little or no irritation.

It should be emphasized that irritation is not a prerequisite for depigmentation; it does, however, accelerate the process. When depigmentation was complete, applications were reduced to once daily. The sites stayed depigmented as long as treatment continued. We observed at least a dozen individuals who were treated daily without mishap for about six months. The only visible skin change was loss of pigment. Depigmentation did not spread beyond the borders of the treated areas.

Repigmentation started within one to two weeks after treatment ceased. This followed the course so well known in vitiligo—inward spread from the borders and centrifugally from the follicular orifices. The pigment was not only restored by about a month but sometimes was increased in relation to normal skin. Generally there was sufficient color blending by three to four months after cessation of treatment to render the treated sites invisible.

Effect of Ultraviolet Radiation

The biweekly application of two minimum erythema doses (MEDs) (determined individually on normal skin) during the period of application was extremely antagonistic to the depigmenting effect. The most that could be achieved was moderate lightening after two months of use. The application of two MEDs of erythemic radiation in depigmented subjects immediately after stopping treatment, with one additional exposure five days later, resulted in complete repigmentation by 10 to 14 days. Irradiated depigmented sites always become hyperpigmented with relation to untreated sites.

Occlusive vs Open Applications

Application twice daily to one side of the back of four individuals was compared to once daily application under a continuous occlusive seal of polyethylene film.

Occlusion increased irritation in

each case. It also enhanced the onset and rate of depigmentation. In three of these four subjects hypopigmentation was evident in a week and was maximal by three to four weeks after the start of treatment.

Variations in the Concentration of Components

The concentration of one ingredient at a time was varied in a group of five subjects.

Hydroquinone.—A hydroquinone concentration of 10% brought about depigmentation more rapidly than that of 5%. Irritancy increased and the cream was more likely to undergo brownish discoloration. This had been noted by Spencer too.¹⁹

Reducing the concentration of the drug to 2% decreased its irritativeness but diminished its potency. Complete depigmentation could be achieved but at the expense of time, maximal effects requiring three months or so. Applications thrice daily almost overcame this.

Tretinoin.—Increasing the concentration of tretinoin to 0.2% accentuated irritation without a corresponding gain in effectiveness. With a concentration of 0.05%, irritancy decreased but a few weeks longer were required for complete depigmentation. Depigmentation was not achieved when the concentration was reduced to 0.01%.

Dexamethasone.—Increasing the concentration of dexamethasone to 0.2% did not lessen irritancy but did enhance effectiveness in a very clear-cut manner. With a 0.05% concentration there appeared to be only a slight loss of effectiveness.

In sum, lowering the concentration of each component by about one half decreases potency, but depigmentation is still obtainable.

Omission of One Ingredient

Omitting one component always results in loss of therapeutic effectiveness; it was not possible to obtain complete depigmentation after three months of twice daily applications in groups of three to four subjects. Lightening was visible but, on the whole, slight. While every ingredient was crucial to attain our established end-point, we could form a crude estimation of the relative

sults

sides, Kahn had come to the conclusion that practically any phenolic substance could induce hypopigmentation with sufficiently intense exposure.¹⁴ He even observed loss of pigment with occlusively applied hexachlorophene.

We tested this possibility by substituting the following substances in place of tretinoin in the basic formula: (1) 2.5% to 5.0% sodium lauryl sulfate (this denatures the horny layer barrier and greatly potentiates penetration); (2) 5% to 10% phenol; (3) 10% resorcinol; (4) 5.0% trichloroacetic acid; (5) 2.5% benzalkonium chloride 3500 (Hyamine 3500) (a cationic quaternary salt); (6) 5% sodium salt of tetrachlorophenol; and (7) 5% salicylic acid.

These studies were conducted on different groups of four to five subjects and always the basic formula was used as a control.

The irritant reactions produced by this group of substances was as great, and in some instances greater, than that produced by the basic formula. In no case, however, was the depigmenting effect comparable. Sodium lauryl sulfate came closest to being an effective replacement, that is to say, the depigmentation was greater than could be achieved with the combination of hydroquinone and the corticosteroid. Phenol was also partially effective. Neither compound can be given serious consideration. Trichloroacetic acid, the quaternary salt, and tetrachlorophenol seemed to nullify the depigmenting effect.

Racial Influence

At first, light- and dark-skinned blacks were used indiscriminately. We gradually came to appreciate that deeply pigmented individuals were more susceptible to the depigmenting effect.

This finding was evaluated in a parallel study in which twice daily applications of the basic formula were made for eight weeks in three groups consisting of five subjects each: (1) very dark blacks, (2) lightly pigmented blacks, and (3) whites of average complexion.

In regard to the onset and final effect after eight weeks of treatment,

there were important

group. Again normal skin was only slightly lightened.

More difficulty was experienced with color blending in patients with postinflammatory hyperpigmentation. These patients were blacks in whom patch tests with irritants and allergens had provoked intensified pigmentation, usually on the forearm or back. Without exception, pigment-loss was first noted in the hyperpigmented square and was regularly well under way by the third

(Fig 4). In our estimation, hydroquinone—no matter how formulated—is not effective enough to satisfy the needs of most patients with hyperpigmentation disorders. Claims such as Spencer's are utterly baffling to us and completely alien to our experience.¹⁸ He achieved depigmentation of senile lentigines on the backs of the hands of older white men in 28 of 41 patients by applying hydroquinone. We found those particular lesions quite resistant.

cytotoxic or at least cytostatic for the epidermis; they decrease epidermal turnover and eventually produce atrophy.²³ Perhaps the secretory product of the melanocyte is suppressed in the same fashion as collagen synthesis is suppressed by fibroblasts.²⁴

We studied specimens of black skin in which epidermal atrophy had been induced by prolonged topical application of a corticosteroid. There was only slight hypopigmentation at the time of biopsy. The quantity of pig-

week of treatment. Needless to say, in this group one seeks only to reduce pigmentation to the background level.

Two individuals with postinflammatory hyperpigmentation responded only moderately; their cases were highly instructive and revealed a limitation that cannot be overcome. In both patients, the skin had become

Spencer also found that 2% hydroquinone would depigment the normal skin of light-skinned blacks in about 50% of the cases. We would emphasize that our inability to achieve anything like that result with the available commercial formulations or with extemporaneous preparations containing 2% to 5% hydroquinone cannot be ascribed to inadequate usage. Our

ment did not seem to be altered. With the dopa stain, however, melanocytes were fewer and often shrunken, with short attenuated dendrites. The intensity of the stain was also diminished. One would normally expect such changes to be accompanied by clinically apparent pigment loss. The probable explanation for this paradox is decreased epidermal turnover. Ko-

intensely hyperpigmented following intensive allergic contact dermatitis. The reason for resistance was seen in

subjects were largely institutionalized volunteers, applications were two to three times daily, and the test

melanocytes move outward at a slowed rate and can thus acquire pigment over a longer time. Production is less

a few days, speeding up the loss of pigment.

It is important to note that there is some balancing-out of the individual effects when the three agents are combined. Clearly, tretinoin overrides the atrophy-producing and antimitotic effect of the corticosteroid. The

use of other potentially irritating substances; washing with soap should be limited to one or two times daily.

Hyperpigmentation rarely begins before three weeks. Do not promise rapid results and urge patience. If hypopigmentation is not under way by two months, the case may be regarded as a treatment failure.

After the desired degree of lightening is obtained, it may be possible to maintain the effect by treatment with low-strength preparations. We are encouraged to think that in some conditions 5% hydroquinone alone will sustain the effect.

Finally, we perforce must mention one potentially nightmarish outcome

of this work, namely, the use of this formula to lighten the skin of normal blacks. We fervently pray that improving social relationships will restrain any dignified black person from that demoralizing practice. Depigmenting normal skin is ethically acceptable only in extensive vitiligo.

REFERENCES

1. Oettel H: Die Hydrochinonvergiftung. Arch Exp Pathol Pharmacol 183:319-362, 1936.
2. Chavin W, Schlesinger W: Some potent melanin depigmentary agents in the black goldfish. Naturwissenschaften 53:413-420, 1966.
3. Denton CR, Lerner AB, Fitzpatrick TB: The inhibition of melanin formation by chemical agents. J Invest Dermatol 18:119-135, 1952.
4. Spencer MC: Hydroquinone bleaching. Arch Dermatol 84:131-134, 1961.
5. Oliver EA, Schwartz L, Warren LH: Occupational leucoderma. Arch Dermatol 42:993-1014, 1940.
6. Lerner AB, Fitzpatrick TB: Treatment of melanin hyperpigmentation. JAMA 152:577-582, 1953.
7. Becker Jr JW, Spencer MC: Evaluation of monobenzone. JAMA 180:279-284, 1962.
8. Dorsey CS: Dermatitis and pigmentary reactions to monobenzyl ether of hydroquinone. Arch Dermatol 81:245-248, 1960.
9. Editorial Comment. Trans St Johns Hosp Dermatol Soc 56:181, 1970.
10. Brun R: Contribution de l'experience de la depigmentation experimentale. Bull Inst Natl Genevoux 61:1-18, 1961.
11. Pathak MA, Frenk E, Szabo G, et al: Cuta-

neous depigmentation. Clin Res 14:272-278, 1966.
12. Bleeher SS, Pathak MA, Hori Y, et al: Depigmentation of skin with 4-isopropyl catachol, mercaptoamines and other compounds. J Invest Dermatol 50:103-110, 1968.
13. Gellin G, Possick PA, Perone VB: Depigmentation from 4-tertiary butyl catachol: An experimental study. J Invest Dermatol 55:190-197, 1970.
14. Kahn G: Depigmentation caused by phenolic detergent germicides. Arch Dermatol 102:177-181, 1970.
15. Malten KE, Seutter E, Hara I, et al: Occupational vitiligo due to paratertiary butylphenol and homologues. Trans St Johns Hosp Dermatol Soc 57:115-131, 1971.
16. Riley PA: The mechanism of hydroxyanisole depigmentation. J Pathol 101:163-167, 1970.
17. Riley PA: Hydroxyanisole depigmentation: In vitro studies. J Pathol 97:193-198, 1969.
18. Spencer MC: Topical use of hydroquinone for depigmentation. JAMA 194:902-904, 1970.